
**EXHIBIT 2** 



TOSO

In this section



ion Exchange (IEC)

Hydrophobic Interaction (HIC)

Affinity (AFC)

#### Size Exclusion (SEC)

Toyopearl HW-40 Toyopearl HW-50 Toyopearl HW-65 Toyopearl HW-65 Toyopearl HW-75 Home What's New Products Literature/Applications Service/Support About Us

Tosoh Bioscience LLC: The Chemistry of Innovation

## Process Media Products

Size Exclusion Chromatography

Toyopearl HW and TSK-GEL PW size exclusion resins are the foundation for all of Tosoh's ion exchange, hydrophobic interaction, and affinity process media.

After an initial polymerization reaction to create the designated particle size and pore size, the resin surface is chemically modified to create the desired final product.

## Toyopearl HW Size Exclusion Beads

Toyopearl HW products are hydroxylated methacrylic polymer resins. Toyopearl HW beads are made commercially in many different pore sizes and particle diameters (30 micron and larger).

They vary from TSK-GEL PW beads by having a lower degree of crosslinking. Lower crosslinking makes available a larger number of resin sites for ligand immobilization when producing Toyopearl HW based ion exchange and hydrophobic interaction chromatography resins.

Therefore a functionalized Toyopearl HW resin will have a higher dynamic binding capacity than the corresponding TSK-GEL PW material.

Because similarly functionalized HW and PW resin types have the same backbone polymer chemistry, the selectivity for proteins and impurities remains the same.

Toyopearl HW based resin products can be used at high linear velocities and withstand operating pressures up to 3 bar.

Resin Chemistry of Toyopearl and TSK-GEL PW resins (Hydroxylated Acrylic)

Note: R = Hydroxylated Aliphatic Group

Comparison of Toyopeari HW resins and TSK-GEL PW resins offered by Tosoh Bioscience

Pore síze(Å)	50	125	400-500	750	1000	>1000	>1700
							November 1884 (Days)
Toyopearl HW:	40	. 50	55	50		75	80
TSK-GEL PW:	1000	2000	4000		5000	6000	
	( miny landadan belands ver	or or a substructive of the Company	tectoroconomicamochecromologichy, phylogene, monec	A COMMON TO THE PROPERTY OF CHEST AND A CONTRACT AN	NO ANTIFECTS STATES TO STATES AND	h month of the control of the contro	O DESCRIPTION OF THE PROPERTY.

Increasing pore surface area

### TSK-GEL PW High Performance Size Exclusion Beads

TSK-GEL PW products are hydroxylated methacrylic polymer resins. TSK-GEL PW beads are made commercially in many different pore sizes and particle diameters (30 micron and smaller).

Media by base resin

They vary from Toyopearl HW beads by having a higher degree of crosslinking (necessitated by the higher flow pressures needed when using smaller particles for chromatography). This decreases the number of resin sites available for ligand immobilization when producing TSK-GE PW ion exchange and hydrophobic interaction chromatography resins.

A functionalized TSK-GEL PW resin will have a lower dynamic binding capacity than the corresponding Toyopearl HW material.

However, chromatographic efficiency, peak width, and resolution are improved because of their smaller particle sizes. In this way they are higher performing chromatography resins than the HV based materials.

Because similarly functionalized HW and PW resin types have the same backbone polymer chemistry, the elution selectivity for proteins and impurities remains the same.

TSK-GEL PW based resin products can be used at high linear velocities and withstand operating pressures up to 20 bar.

Tosoh Bioscience LLC 156 Keystone Drive, Montgomeryville, PA, 18936 Tel: (215)



# **TOSOH BIOSCIENCE LLC**

156 Keystone Drive, Montgomeryville, PA 18936

Orders and Services: 800-366-4875

## Toyopearl HW-65

**Toyopearl HW-65** is a hydroxylated methacrylic polymer bead for size exclusion chromatography of proteins between 40,000-5,000,000 Da. It also is the base bead for all Toyopearl "-650" designated products.

#### Structure:

Note: R = Hydroxylated Aliphatic Group

### **Product Attributes:**

Pore size (mean): 1,000Å

Exclusion limit: 5,000,000 Da

Particle pize (mass): 45µm (F-grade)

Particle size (mean): 45µm (r-grade) 30µm (S-grade)

Pressure rating: 3 bar


Shipping buffer 20% ethanol

pH stability: 2-13

shelf life: 10 years

Comparative information for SEC resins

# Calibration curves for globular proteins on Toyopearl HW resins



Coloran: 22mmiO x 30cm Sample: protein steedards


Elution: 0.06M phosphate buffer (pH 7.0) in 0.06M KCi Legend: V = elution volume V = column volume

Typical resolution changes based on particle size

Toyopearl HW-65 Page 2 of 2

### Literature:

Size Exclusion Chromatography



Column

Toyopeari HW-55, 26mmlD x Nicm 33.3mM phosphate buffer (pH7.0), 0.2M NaCl Elegant:

Flow rate: 106mL/h (20cm/h)

Temp.: 250 Cetector: UV #280mm

1. Thyroglobusin (0.3%), 2. + Globusin (0.3%) Sample:

3. (i-Lactoglobulin (0.3%), 4. Oyoochromin C (0.1%)

ing Vol.: 1ml

## Ordering information:

(For quantities up to 1L please download our Price List; for pricing above 1L or to place your order, please contact customer service)

PIN	Description	Resin volume	Particle size
19814	Toyopearl HW-65F	150mL	30-60µm
07465	Toyopearl HW-65F	500mL	30-60µm
14689	Toyopearl HW-65F	1L	30-60µm
08069	Toyopearl HW-65F	5L	30-60µm
19815	Toyopearl HW-65S	150mL	20-40µm
07467	Toyopearl HW-65S	250mL	20-40µm
07464	Toyopearl HW-65S	500mL	20-40µm
14688	Toyopearl HW-65S	1L	20-40µm
08068	Toyopearl HW-65S	5L	20-40µm

**EXHIBIT 3** 



# TOSOH BIOSCIENCE LLC

156 Keystone Drive, Montgomeryville, PA 18936

Orders and Services: 800-366-4875

## **Toyopearl AF-Amino-650**

**Toyopearl AF-Amino-650** is a reactive resin used for the coupling of specific ligands for affinity chromatography. Ligands are immobilized by either peptide bond formation or reductive amination through their respective carboxylate or aldehyde groups.

#### Structure:

(HW-65) - O-R-O-CH<sub>2</sub>-CH-CH<sub>2</sub>-NH<sub>2</sub>

Approximate Ligand Density: 100µmol/mi.

## Coupling procedure

(FW-95)-0-R-0-CH<sub>2</sub>-CHOH-CH<sub>2</sub>NH<sub>2</sub> + HOOC-Ligand

Toyopean Af-Amino carinelimite this &

(FIVE)-O-R-O-CH2-CHOH-CH3NH-CO-Ligand

Peptide bood linkage

#### **Product Attributes:**

Pore size (mean): 1,000Å

Particle size (mean): 65µm (M-grade)

Pressure rating: 3 bar

Shipping buffer: 20% ethanol

pH stability: 2-13 Shelf life: 10 years (HW-65)-O-R-O-CH<sub>2</sub>-CHOH-CH<sub>2</sub>NH<sub>2</sub> + OHC-Ligand
Topopeani AF-Assure | PH-7
25°C

(HW-65)-O-R-O-CH<sub>2</sub>-CHOH-CH<sub>2</sub>-N=CH-Ligand
Schill's Base Intermediate
| NaCNBH<sub>2</sub>

(HW-65)-O-R-O-CH<sub>2</sub>-CHOH-CH<sub>2</sub>-NH-CH<sub>2</sub>-Ligand
Stable secondary amins linkage

## Literature:

## Affinity Chromatography

## Ordering information:

(For quantities up to 1L please download our <u>Price List</u>; for pricing above 1L or to place your order, please contact <u>customer service</u>)

R » hydrophilic

PiN	Description	Resin volume	Ligand density
43411	Toyopearl AF-Amino-650M	10mL	70 - 130µeq/mL
08002	Toyopearl AF-Amino-650M	25mL	70 - 130µeq/mL
08039	Toyopearl AF-Amino-650M	100mL	70 - 130µeg/mL

18074	Toyopearl AF-Amino-650M	1L	70 - 130µeq/mL
18316	Toyopearl AF-Amino-650M	5L	70 - 130µeq/mL